

SURVEY OF LIVESTOCK INFLUENCES ON STREAM AND RIPARIAN ECOSYSTEMS IN THE WESTERN UNITED STATES

(Published by *Journal of Soil and Water Conservation*, 1999, Vol. 54, pp. 419-431)

A.J. Belsky*, A. Matzke#, S. Uselman#

*Staff Ecologist, #Research Associates Oregon Natural Desert Association, 732 SW 3rd Ave., Suite 407, Portland OR 97204,

Telephone/Fax 503-228-9720, email jbelsky@onda.org

Acknowledgments: We thank R. Amundson, H. Campbell, D. DeLong, T. Dudley, D. Ferguson, N. Ferguson, S. Fouty, A. Kerr, T. Myers, M. O'Brien, E. Painter, R. Phillips, W. Platts, and J. Rhodes for providing insightful comments on drafts of this paper, as well as Northwest Fund for the Environment and the Bullitt Foundation for financial support.

Interpretive Summary

Livestock grazing has damaged approximately 80% of stream and riparian ecosystems in the western United States. Although these areas compose only 0.5-1.0% of the overall landscape, a disproportionately large percentage (~70-80%) of all desert, shrub, and grassland plants and animals depend on them. The introduction of livestock into these areas 100-200 years ago caused a disturbance with many ripple effects. Livestock seek out water, succulent forage, and shade in riparian areas, leading to trampling and overgrazing of streambanks, soil erosion, loss of streambank stability, declining water quality, and drier, hotter conditions. These changes have reduced habitat for riparian plant species, cold-water fish, and wildlife, thereby causing many native species to decline in number or go locally extinct. Such modifications can lead to large-scale changes in adjacent and downstream ecosystems.

Despite these disturbances, some people support continued grazing. These advocates argue that most of the damage occurred 50-100 years ago; however, recent studies clearly document that livestock continue to degrade western streams and rivers, and that riparian recovery is contingent upon total rest from grazing.

Abstract

This paper summarizes the major effects of livestock grazing on stream and riparian ecosystems in the arid West. We focused primarily on results from peer-reviewed, experimental studies, and secondarily on comparative studies of grazed vs. naturally or historically protected areas. Results were summarized in tabular form.

Livestock grazing was found to negatively affect water quality and seasonal quantity, stream channel morphology, hydrology, riparian zone soils, instream and streambank vegetation, and aquatic and riparian wildlife. No positive environmental impacts were found. Livestock were also found to cause negative impacts at the landscape and regional levels. Although it is sometimes difficult to draw generalizations from the many studies, due in part to differences in methodology and environmental variability among study sites, most recent scientific studies document that livestock grazing continues to be detrimental to stream and riparian ecosystems in the West.

Introduction

Grazing by livestock has damaged 80% of the streams and riparian ecosystems in arid regions of the western United States (U.S. Department of the Interior (USDI) 1994a). A number of symposia (e.g. Warner and Hendrix 1984, Johnson et al. 1985, Gresswell et al. 1989, Meehan 1991, Clary et al. 1992) and reviews (Platts 1981b, 1982, 1991, Kauffman and Krueger 1984, Skovlin 1984, Chaney et al. 1990, 1993, Armour et al. 1994, Fleischner 1994, Rhodes et al. 1994, USDI 1994a, Kattelman and Embury 1996, Ohmart 1996) describe this degradation. Livestock grazing affects watershed hydrology, stream channel morphology, soils, vegetation, wildlife, fish and other riparian-dependent species, and water quality at both local and landscape scales. Because riparian and stream ecosystems represent only 0.5-1% of the surface area of arid lands of the eleven western United States (U.S. General Accounting Office (US-GAO) 1988, Chaney et al. 1990, Ohmart 1996), they were historically ignored by land managers. In fact, riparian habitats in the West were viewed until the late 1960s as "sacrifice" areas (e.g., Stoddart and Smith 1955), being dedicated primarily to providing food and water for domestic livestock.

Recently, both critics and advocates of arid-land livestock grazing have focused their attention on western streams and their associated riparian zones, especially those in shrublands, grasslands, and deserts of the Southwest, Great Basin, and Pacific Northwest. Critics of grazing emphasize damage to riparian habitats to illustrate the unsuitability of cattle grazing in the arid West, while advocates of grazing argue that most of the damage to land and streams occurred 50-100 years ago, before modern grazing systems were instituted.

The evidence is undeniable that early grazing practices -- before the Taylor Grazing Act in 1934 established some control over livestock grazing in the public domain -- were highly destructive (Duce 1918, Bryan 1925, Leopold 1946). However, recent studies document that livestock grazing remains a key factor in the continued degradation of riparian habitats (US-GAO 1988, Szaro 1989, Platts 1991, Elmore and Kauffman 1994, Fleischner 1994, McIntosh et al., 1994, USDI 1994a, Ohmart 1996). As recently as 1990, Chaney et al. (1990) wrote in a U.S. Environmental Protection Agency (US-EPA) report on livestock grazing that "extensive field observations in the late 1980s suggest that riparian areas throughout much of the West were in their worst condition in history" (p.5). A joint Bureau of Land Management (BLM) and US Forest Service Report (USDI 1994a)

also concludes that "riparian areas have continued to decline [since 1934]" (p.25) and estimates that 20% of the riparian areas managed by BLM are "non-functioning" and 46% are "functioning at risk." Altogether, less than 20% of potential riparian habitat in the western United States still exists (USDI 1994a). This continued decline has been attributed, in part, to increased numbers of cattle in western rangelands (Trimble and Mendel 1995); between 1940 and 1990, the number of cattle in the western United States increased from 25,500,000 to 54,400,000.

Recent scrutiny by scientists reflects a growing recognition by the public, land managers, and scientific community of the importance of streams, rivers, and riparian habitats to western ecosystems. One reason for this interest is the high productivity and biodiversity of riparian systems, which is due, in part, to their high soil moisture and fertility levels (Hubbard 1977, Meehan et al. 1977, Thomas et al. 1979, Knight and Bottorff 1984, Fleischner 1994, Ohmart 1996). Riparian areas in arid and semi-arid regions are composed of complex edaphic and vegetation mosaics because of high variability in landforms, soil types, and location of surface and subsurface water (Thomas et al. 1979, Green and Kauffman 1995, Lee et al. 1989, Gregory et al. 1991). These mosaics, plus extensive borders (ecotones) between moist streambanks and arid uplands, result in high species diversity (Thomas et al. 1979, Lee et al. 1989). An estimated 60-70% of western bird species (Ohmart 1996) and as many as 80% of wildlife species in Arizona and New Mexico (Chaney et al. 1990) and in southeastern Oregon (Thomas et al. 1979) are dependent on riparian habitats. Consequently, riparian ecosystems are considered to be important repositories for biodiversity throughout the West.

Riparian zones provide key services for all ecosystems, but are especially important in dry regions, where they provide the main source of moisture for plants and wildlife, and the main source of water for downstream plant, animal, and human communities (Meehan et al. 1977, Thurow 1991, Armour et al. 1994, among others). These services are highly dependent on streambanks and flood plains being in a vegetated and relatively undisturbed state. Rooted streamside plants retard streambank erosion, filter sediments out of the water, build up and stabilize streambanks and streambeds, and provide shade, food, and nutrients for aquatic and riparian species (Winegar 1977, Thomas et al. 1979, Kauffman and Krueger 1984). The ability of undisturbed plant communities to stabilize banks was notable during extensive floods in eastern Oregon in 1996, when shrubby vegetation in ungrazed sections of the Deschutes River "broke the flood's velocity and combed logs and mud from the river" (Meehan 1996).

Healthy riparian areas also act as giant sponges during flood events, raising water tables and maintaining a source of streamwater during dry seasons. The result is a more stable streamflow throughout the year (US-GAO 1988).

Cattle cause more damage to riparian zones than their often small numbers would suggest. Cattle tend to avoid hot, dry environments and congregate in wet areas for water and forage, which is more succulent and abundant than in uplands. They are also attracted to the shade and lower temperatures near streams, most likely because their species evolved in cool, wet meadows of northern Europe and Asia. In fact, cattle spend 5-30

times as much time in these cool, productive zones than would be predicted from surface area alone (Roath and Krueger 1982, Skovlin 1984). One study found that a riparian zone in eastern Oregon comprised only 1.9% of the grazing allotment by area, but produced 21% of the available forage and 81% of forage consumed by cattle (Roath and Krueger 1982).

Our goal is to summarize along biological and ecological lines the major effects of cattle grazing in stream and riparian ecosystems. We include only those studies that discuss the direct and indirect effects of livestock activities on stream and riparian habitats. We exclude other aspects of livestock production such as conversion of flood plains to cultivated fields for livestock feed, leaching of fertilizer from these fields into streams, and streamwater diversion for crop or pasture irrigation. We also do not include the effects of impounding streamwater for stock ponds or other activities that support livestock production, although these activities contribute significantly to stream degradation.

Methods

We searched the scientific literature for peer-reviewed empirical papers and reviews of the biological and physical effects of livestock on western rivers, streams, and associated riparian areas. Because of the extensive literature on the subject, not all papers could be reviewed or cited. In choosing the papers to be included, we gave highest priority to recent papers in refereed journals presenting experimental manipulations such as paired samples from grazed vs. ungrazed areas or from heavily grazed vs. lightly grazed pastures (when ungrazed controls were not included in the experimental design). Many of these studies used sites recently protected from grazing as controls (e.g., Kauffman et al. 1983a, Schulz and Leininger 1990), but a few used previously ungrazed areas to which livestock were newly introduced (e.g., Sedgwick and Knopf 1987, Samson et al. 1988). Secondary priority was given to descriptive or comparative studies of grazed vs. naturally or historically protected areas where similarity of initial conditions could be inferred. Where there was a paucity of data, we also used non-peer-reviewed reports, usually from government documents or symposia. In no case were our general conclusions drawn from unrefereed reports or from studies showing anomalous results. Instead, we based our conclusions on what seemed to be the consensus of experts in the field.

We also identified and listed comprehensive review papers on each topic. Environmental impacts were defined as environmental changes that were significant at the $P < 0.1$ level (e.g., Peterman 1990) (discussed below) or those effects deemed significant by the authors.

Results

Damage caused by cattle to riparian and stream habitats in the arid and semi-arid West can be separated into two broad categories: impacts that occur at the local level (Table 1) and those that occur at landscape and regional levels (Table 2). Local impacts can be further segregated by their effects on water quality and seasonal quantity, stream channel morphology, hydrology, riparian-zone soils, instream and streambank vegetation,

aquatic biota, and terrestrial wildlife (Table 1). Local impacts have been investigated in a large number of studies, but landscape-level impacts have received less attention.

Our search uncovered no systematic investigations showing positive impacts or ecological benefits that could be attributed to livestock activities when grazed areas were compared to protected areas (see also Bock et al. 1993, Ohmart 1996). Thus, we mostly present negative environmental impacts. In general, there was little debate about the effects of livestock grazing. Most authors tended to agree that livestock damage stream and riparian ecosystems.

Discussion

In the following, we discuss pertinent topics that have not been addressed in depth in recent reviews. These reviews, which are listed after each major category in Tables 1 and 2, should be consulted for additional discussion of other topics.

Positive and neutral effects of cattle grazing on riparian zones. An extensive literature search did not locate peer-reviewed, empirical papers reporting a positive impact of cattle on riparian areas when those areas were compared to ungrazed controls, but some studies reported no statistically significant effects due to riparian grazing (e.g., Buckhouse and Gifford 1976, Samson et al. 1988). The authors of these papers usually explained this absence of statistically significant impacts as being due to stochastic or design problems associated with their research, rather than to grazing having no effect on vegetation, fish, or stream hydrology. They described such problems as (1) high variability among treatment plots, which masked treatment effects (e.g., Tiedemann and Higgins 1989, Shaw 1992), (2) insufficient recovery periods after protection from grazing (e.g., Hubert et al. 1985, Sedgwick and Knopf 1991, Shaw 1992, Sarr et al. 1996), (3) heavy browsing and grazing by native herbivores (or trespassing cattle) on supposedly ungrazed control plots (e.g., Shaw 1992, Clary et al. 1996), (4) unplanned disturbances such as flooding (e.g., Sedgwick and Knopf 1991, Clary et al. 1996, Myers and Swanson 1996a), and (5) the unknown effects of a prior history of heavy grazing, which may have permanently altered stream function and prevented recovery of control plots (e.g., Tiedemann and Higgins 1989).

The absence of significant effects may also be due to investigators setting statistical significance at arbitrarily low levels (i.e., at $P < 0.05$). Peterman (1990) argues that many studies, such as those with few treatment replications or high spatial variability, have low power (i.e. poor ability) to detect environmental change. Because of the possibility that already depleted fish stocks could become endangered or important habitats become permanently altered, he argues that higher probability levels (i.e., $P < 0.1$) are appropriate to test significance of hypotheses.

Authors have also attributed non-significant results to supplemental feeding of livestock (e.g., Sedgwick and Knopf 1991), which resulted in lower forage consumption levels than originally prescribed, and to high recreational fishing, which obscured the negative effects of grazing on fish populations (e.g., Hubert et al. 1985). Finally, severe

environmental damage such as loss of native species or channel downcutting cannot be reversed in just a few years of protection. Streams may recover slowly or only over geological time scales (Sarr et al. 1996). Together, these circumstances have caused some (e.g., Platts 1982) to question the ability of many experimental techniques to adequately assess livestock impacts. Others (e.g. Peterman 1990) also question the statistical power of many experiments to accept or reject hypotheses.

Several recent papers (e.g., Clary and Webster 1989, Elmore and Kauffman 1994, Burton and Kozel 1996, Weller 1996) describe the benefits of reduced cattle stocking rates and newer grazing systems, such as seasonal grazing, rest-rotation, and deferred grazing. The authors also discuss examples of grazed riparian zones regaining their herbaceous and woody cover and water quality. These studies, however, only contrasted newer grazing systems with more traditional and destructive systems, such as year-long grazing and high stocking rates. They did not contrast these systems with no-grazing. The only conclusion that could be fairly drawn from these studies is that newer grazing systems improve streamside conditions relative to other grazing systems, not that cattle grazing truly benefit riparian zones. In fact, Meehan and Platts (1978) and Platts and Wagstaff (1984) found no grazing system that was compatible with healthy aquatic ecosystems.

In mid-western prairies, livestock have been reported to be useful at breaking up dense, rank vegetation near wetlands (Weller 1996). However, in the Intermountain West, where low densities of native grazers provided only light grazing and trampling disturbances during the last 10,000 years, riparian species have inherently lower tolerances for livestock disturbances (Mack and Thompson 1982). It is doubtful that grazing or trampling by cattle in this region would do more good than harm.

Problems in drawing generalizations from riparian studies. Although most research has shown grazing in streams and riparian zones to be deleterious, results have been variable (Platts 1982, Trimble and Mendel 1995). This has caused riparian specialists problems in drawing broad generalizations about the effects of cattle grazing. These problems can be attributed to several issues:

1. *Inadequacy of study design.* Most watershed-scale riparian management plans were not designed as experiments with the idea of researchers evaluating them years later.
2. *Inherent variability found between and within watersheds.* Streams are unique, having their own combination of channel morphology, soils, climate, riparian species, geology, and hydrology (Elmore and Beschta 1987, Myers and Swanson 1991, Trimble and Mendel 1995). One management strategy may have a particular effect in one area, but a greater or lesser effect elsewhere.
3. *Insufficient study replication.* Lack of adequate replication of experimental treatments make data interpretation difficult (Matthews 1996).

4. *Ambiguities or differences in study design* (Platts 1982, Rinne 1985). In some cases, terms such as "heavy" and "light" grazing to describe grazing treatments are subjective, making comparisons within and between experimental studies difficult (Fleischner 1994, Trimble and Mendel 1995). In other cases, differences in research methodologies make comparisons unreliable (Trimble and Mendel 1995).
5. *Grazing inside exclosures by small mammals and invertebrates*. Small animals often congregate inside exclosures where food and cover are abundant. Increases in grazing inside exclosures by grasshoppers, rabbits, and rodents may reduce differences between treatments, thus masking the effects of cattle grazing outside the exclosure.
6. *Prior grazing history*. Many pastures now protected within exclosures were grazed at some time in the past and thus do not accurately fall within a truly ungrazed (i.e. "pristine") landscape. In fact, many older exclosures were purposely erected in severely overgrazed and eroded areas in order for investigators to monitor recovery and successional processes. Since many of these protected stream segments may have been deeply downcut previously, their recovery may take hundreds to thousands of years. These exclosure studies, therefore, may underestimate the true extent of livestock damage because they fail to take into account the damage that occurred before the exclosures were erected (Fleischner 1994).
7. *Variable time lags*. Recovery of different ecological, hydrological, and geomorphologic processes require different amounts of time, often longer than the average research grant and sometimes longer than the life-span of the researcher. Recovery of herbaceous and woody vegetation along stream sides may begin immediately after grazing is terminated, while the recovery of channel form may take hundreds of years (Kattelman and Embury 1996, Trimble and Mendel 1995, Clary et al. 1996).
8. *Influences from outside the study area*. Stream channel morphology and aquatic organisms respond not only to factors occurring inside the study area, but to those occurring outside as well (Rinne 1985). Soil compaction and reduced infiltration of rainwater due to cattle trampling on slopes above riparian exclosures may increase the volume of water flowing over soil surfaces and into protected research sites. In addition, grazed streambanks upstream from exclosures may fail, releasing sediments into protected segments. Together, these factors may contribute large amounts of sediment to the stream system, inhibiting stream recovery (Kondolf 1993). Similarly, water flowing out of exclosures may be cleaner, cooler, and produce better spawning habitat downstream than that inside the exclosures (Duff 1977, Rinne 1985). Conditions over the larger landscape, therefore, minimize differences in grazed/ungrazed comparative studies.

In spite of numerous problems in experimental design and difficulties in interpreting earlier studies, Platts (1982) concluded that livestock grazing was the major cause of degraded stream and riparian environments and reduced fish populations throughout the arid West. In an extensive review of the literature, he found that 85% of

the studies demonstrated that livestock negatively impacted riparian and stream ecosystems, which he concluded was a sufficiently powerful statistic to override inadequacies in individual experimental design.

Effects of riparian grazing on channel morphology and water tables. Plants on undisturbed uplands and streambanks slow the downhill flow of rainwater, promoting its infiltration into soils. Water that percolates into the ground moves downhill through the sub-soil and seeps into stream channels throughout the year, creating perennial flows. But as upland and riparian vegetation is removed by livestock and as hillsides and streambanks are compacted by their hooves, less rainwater enters the soil and more flows overland into streams, creating larger peak flows. This was illustrated in a simulation by Trimble and Mendel (1995), who estimated that peak storm runoff from a 120 ha basin in Arizona would be 2-3 times greater when "heavily" grazed than when "lightly" grazed. Moderate and high rainfall events in grazed sites are, therefore, more likely to result in high energy and erosive floods, which deepen and reshape stream channels (Fig. 1, USDI 1994a).

Where streams flow over deep soils or unconsolidated substrates, the erosive energy of floods cause channel downcutting, or incision (Fig. 1). As the channel deepens, water drains from the flood plain into the channel, causing a lowering (subsidence) of the water table. The roots of riparian plants are left suspended in drier soils. Eventually, riparian plants and their associated wildlife species are replaced by upland species such as sagebrush (*Artemisia* spp.) and juniper (*Juniperus* spp.), which can tolerate these drier soils. Additionally, with less water entering upslope and riparian soils, less is available to provide late-season flows. Consequently, the high intensity floods of the spring and early summer are often followed by low and no flow in late summer and fall.

Effects of riparian grazing on biodiversity. Most studies comparing grazed and protected riparian areas show that some plant and animal species decrease in abundance or productivity in grazed sites while other species increase. Plant species that commonly decline with livestock grazing are either damaged by removal of their photosynthetic and reproductive organs, or are unable to tolerate trampling or the drier conditions caused by lowered water tables. Plant species that commonly increase with livestock grazing are usually weedy exotics that benefit from disturbed conditions, upland species that prefer the drier conditions created by grazing, or sub-dominant species that are released from competition when taller neighbors are grazed down (Kauffman and Krueger 1984, Schulz and Leininger 1991, Stacy 1995, Green and Kauffman 1995, Ohmart 1996, Sarr et al. 1996).

Neotropical migratory birds (Bock et al. 1993, Saab et al. 1995) and prairie waterbirds (Weller 1996) are also variously affected by livestock grazing. After reviewing a large number of relevant studies, Saab et al. (1995) concluded that livestock grazing in the West led to a decline in abundance of 46% of the 68 neotropical migrant landbirds that utilize riparian habitat, an increase in 29% of the migrants, and no clear response in 25%. Those species that are grounded nesting or forage in riparian areas with heavy shrub or ground cover tended to decrease in abundance with grazing, while species that prefer open habitats, are ground foragers, or are attracted to livestock (i.e., cowbirds (*Molothrus*

spp.), tended to increase in abundance in grazed riparian habitats (Bock et al. 1993, Saab et al. 1995). Cavity and canopy nesters were least affected. After a thorough analysis, Bock et al. (1993) concluded that few neotropical bird species actually "benefited from [cattle] grazing in riparian habitats, and that those that do are not restricted to riparian communities" (p.302). In other words, species that benefit from grazing are already widely distributed over the landscape and gain no extra benefit from additional habitat. Conversely, those species that are harmed by grazing are usually restricted to riparian habitats. Riparian grazing, therefore, makes them vulnerable to local extinction.

Fish populations are also differentially affected by livestock grazing. As stream waters become warmer and more sediment-laden due to streamside grazing (Table 1), trout, salmon, and other cold-water species decline in number and biomass. They are often replaced by less valued and more tolerant species. For example, Stuber (1985) found a higher biomass of game fish (predominantly brown trout (*Salmo trutta*)) in protected stream segments in Colorado, but a higher biomass of non-game species (predominantly longnose sucker (*Catostomys catastomus*)) in grazed segments. Similarly, Marcuson (1977) found that trout (*Salmo* spp.) were more abundant in an ungrazed stream segment in the Beartooth Mountains while mountain whitefish (*Prosopium williamsoni*) were more abundant in a grazed segment.

Changes in species composition due to cattle grazing should not be evaluated in conventional species-diversity terms, since even an influx of exotic weeds will increase species richness and diversity. These weeds may increase diversity, but they also alter wildlife habitat and ecosystem processes (i.e. erosion rates, seasonal flows) to which native species are adapted. Of greater importance than species diversity is whether grazing reduces the abundance or diversity of native species and riparian specialists, and whether these species are being replaced by introduced or upland species. In both cases, such changes lead to a reduction in native biological diversity, homogenization of the biotic landscape, and loss of high-value wildlife (i.e. game) species (Stuber 1985, Bock et al. 1993). Reductions in number, size, and productivity of native riparian or aquatic species are nearly always viewed as negative or as representing declining ecosystem health (Ohmart 1996).

Cattle grazing has converted many of the riparian habitats in the arid West into communities dominated by habitat generalists and weedy species such as dandelions (*Taraxacum officinale*), cheatgrass (*Bromus tectorum*), cowbirds, and small-mouth bass (*Micropterus dolomieu*), and by upland or abundant species such as sagebrush, juniper, and speckled dace (*Rhinichthys osculus*). As a result, both habitat quality and native species diversity have been severely reduced (Marcuson 1977, US-GAO 1988, Armour et al. 1994, Popolizia et al. 1994, Green and Kauffman 1995, Sarr et al. 1996). Consequently, a recent Forest Service report found livestock grazing to be the fourth major cause of species endangerment in the United States and the second major cause of endangerment of plant species (Flather et al. 1994). Within certain regions (i.e. Arizona Basin and Colorado/Green River Plateau), livestock grazing was listed as the #1 cause of species being federally listed as threatened or endangered.

Effects of riparian grazing on water quality. Bacterial contamination of drinking and surface water by domestic livestock is a significant non-point source of water pollution (George 1996). Although usually not considered pathogenic (Gary et al. 1983), fecal coliform (e.g., *Escherichia coli*), and enterococci bacteria are regularly monitored in surface waters because they are indicators of fecal contamination that may include pathogenic organisms such as *Cryptosporidium*, *Giardia*, *Salmonella*, *Shigella* and enteric viruses (Bohn and Buckhouse 1985b, George 1996). Because these organisms are carried by cattle and because fecal bacteria levels tend to increase with increasing grazing pressure (Gary et al. 1983, Owens et al. 1989, George 1996), the probability of disease-causing organisms contaminating swimming areas and entering human water supplies increases with intensity of cattle use.

Another concern is that nutrients found in animal wastes stimulate algal and aquatic plant growth when they are deposited directly or washed into streamwater. If resulting plant growth is moderate, it may provide a food base for the aquatic community. If excessive, these nutrients stimulate algal blooms. Subsequent decomposition of the algae leads to low dissolved oxygen concentrations (US-EPA 1995), which endangers aquatic organisms.

Landscape and regional effects of riparian grazing. The impacts of grazing on local riparian and stream environments and on stream morphology may be acute, but they also often extend far beyond their immediate surroundings (Table 2). Streams connect uplands to lowlands, terrestrial ecosystems to aquatic, and arid ecosystems to moist (Gregory et al. 1991, Knopf and Samson 1994). They act as corridors for migrating animals, provide moisture for aquatic, riparian, and upland species, and distribute sediments and nutrients downstream (Table 2; Thomas et al. 1979, Lee et al. 1989). In the case of anadromous fish, nutrients that are consumed in the ocean are brought inland, where they are distributed throughout the landscape as the fish are consumed by predators or decompose along streambanks after spawning.

By degrading water supplies and reducing the area of healthy riparian habitat, livestock fragment these landscape-level connections. They also damage the connection between natural and human communities, since degraded streams reduce the potential for recreational fishing and swimming, degrade municipal water supplies, provide less water for reservoirs, and damage coastal commercial fishing.

Neither are streams isolated from their adjacent uplands. Heavy grazing on upland communities impacts riparian areas primarily by increasing runoff and erosion. Blackburn (1984) and Trimble and Mendel (1995) summarized the negative impacts of heavy grazing on watersheds. They listed the erosive force of raindrops on denuded surfaces, the shearing force of hooves on slopes, decreased soil organic matter, and increased soil compaction as primary impacts. Together, these lead to reduced water infiltration and increased runoff, soil bulk density, erosion, and sediment delivery to streams. In addition, cattle form trails and terracettes (Trimble and Mendel 1995) (also called bovine terraces), which are also subject to erosion (Rostagno 1989).

Other factors contributing to riparian degradation. Cattle grazing is not the only factor damaging stream and riparian habitats in the arid West. Urban development, mining, damming for hydroelectric power, road construction, local eradication of beaver, logging, agricultural activities, and water diversions for industry, irrigation, and municipal water supplies have also exacted heavy tolls on riparian and aquatic ecosystems (Skovlin 1984, Szaro 1989, USDI 1994b). These factors acting alone and in combination have caused devastating cumulative impacts on western streams (Lee et al. 1989). Despite this, livestock grazing is still considered to be the most pervasive source of upland and riparian habitat degradation in the arid West (Elmore and Kauffman 1994, USDI 1994a, Ohmart 1996, among others).

Effects of riparian grazing in humid environments. Most investigations of the effects of livestock grazing on streams, rivers, and riparian zones have been located in arid regions. Although empirical studies from more humid (mesic) regions, such as western Oregon and Washington, the mid-West, and the eastern United States, are not as numerous (Trimble and Mendel 1995), available evidence suggests that environmental impacts of grazing in these regions are similar to those in drier areas. In all environments, cattle consume streamside vegetation, disturb soils, destabilize streambanks, deposit manure and urine, and churn up channel sediments (Trimble 1994, Armour et al. 1994, Trimble and Mendel 1995). Similar to arid areas, cattle were found to reduce overhead cover, herbaceous cover on banks, and woody vegetation in western Washington and Wisconsin (Chapman and Knudsen 1980, White and Brynildson 1967). Livestock also increased concentrations of ammonia, nitrate, soluble phosphate, chemical oxygen demand, and total organic carbon in runoff in Nebraska (Schepers and Francis 1982), increased concentrations of organic nitrogen, organic carbon, and sediment in runoff in Ohio (Owens et al. 1989, 1996), caused streambank erosion in Pennsylvania (Davis et al. 1991) and Tennessee (Trimble 1994), and increased soil loss in North Dakota (Hofmann and Ries 1991).

In some cases grazing may be even more damaging in wetter than in drier environments because moist soils are more vulnerable to compaction and disturbance than dry soils (Marlow and Pogacnik 1985, Trimble and Mendel 1995, McInnis 1996). In other cases, damage to riparian and stream habitats may be less severe in wetter climates because cattle may be less attracted to streambanks in areas where upland grasses are green and palatable for more months of the year.

Conclusion

The current debate over the environmental impacts and suitability of livestock grazing in arid western ecosystems has resulted in supporters declaring that livestock sometimes benefit streams (Savory 1988). Nearly all scientific studies, both observational and experimental, refute this claim. Livestock do not benefit stream and riparian communities, water quality, or hydrologic function in any way (Table 1). However, their damage can be reduced by improving grazing methods, herding or fencing cattle away from streams, reducing livestock numbers, or increasing the period of rest from grazing (Armour et al. 1994, Elmore and Kauffman 1994). The conclusion that all grazing

practices detrimentally affect riparian areas (Elmore and Kauffman (1994) is to be expected since traditional grazing systems were developed for protecting upland grasses, not for protecting riparian plants and streambanks (Platts 1991, Saab et al. 1995).

With improved livestock management, previously denuded streambanks may revegetate and erosion may decline (Elmore and Kauffman 1994), but recovery will take longer than if grazing were terminated completely (Myers and Swanson 1995, 1996a, Ohmart 1996). Trimble and Mendel (1995) concluded that "although there may have been improvements in grazing management, the increase of cattle in the West [a doubling over the last 50 years] suggest that grazing impacts will continue into the foreseeable future" (p 233).

New studies go even further by suggesting that new grazing systems have only served to slow the rate of degradation, not reverse it. Sarr et al. (1996), for example, found that ten full years of livestock exclusion was necessary to reverse a negative trend and allow stream conditions to begin to improve. Elmore and Kauffman (1994) best summed up available evidence by stating that "livestock exclusion has consistently resulted in the most dramatic and rapid rates of ecosystem recovery " (p. 216).

Although the possibility of streams recovering their plant cover and ecological functions while providing food and water for livestock use is appealing (i.e. a win-win situation), it is largely contradicted by existing evidence (Table 1). Riparian specialist Robert Ohmart of the University of Arizona questions whether weakened and degraded riparian communities throughout the arid West can "hang onto their thread of existence for another 30-50 years" (Ohmart 1996, p. 272) while waiting for grazed systems to recover.

All discussions of improved grazing systems allude to the fact that the best prescription for stream recovery is a long period of rest from livestock grazing. Even those who strongly believe grazing to be compatible with healthy riparian ecosystems point out that 2-15 years of total livestock exclusion is required to initiate the recovery process (Duff 1977, Skovlin 1984, Clary and Webster 1989, Elmore 1996, Clary et al. 1996). Consequently, streams that are permanently protected from grazing have the highest probability of successful recovery (Claire and Storch 1977, Chaney et al. 1990, Bock et al. 1993, Armour et al. 1994, Fleischner 1994, Rhodes et al. 1994, Ohmart 1996, Case and Kauffman 1997).

Literature cited

Armour, C., D. Duff, and W. Elmore. 1994. The effects of livestock grazing on western riparian and stream ecosystem. *Fisheries* 19(9):9-12.

Atwill, E.R. 1996. Assessing the link between rangeland cattle and water-borne *Cryptosporidium parvum* infection in humans. *Rangelands* 18:48-51.

Belsky, A.J., and D.M. Blumenthal. 1997. Effects of livestock grazing on stand dynamics and soils in upland forests of the interior West. *Cons. Biol.* 11:315-327.

Blackburn, W.H. 1984. Impact of grazing intensity and specialized grazing systems on watershed characteristics and responses. p. 927-983. In: *Developing strategies for range management*. Westview Press, Boulder, CO.

Bock, C.E., V.A. Saab, T.D. Rich, and D.S. Dobkin. 1993. Effects of livestock grazing on neotropical migratory landbirds in western North America. p. 296-309. In: D.M.

Finch, P.W. Stangel (eds.), Status and management of neotropical migratory birds. USDA Forest Serv. Gen. Tech. Rep. RM-229.

- Boggs, K., and T. Weaver. 1992. Response of riparian shrubs to declining water availability. p. 48-51. In: W.P. Clary, E.D. McArthur, D. Bedunah, and C.L. Wambolt (compilers), Proceedings-Symposium on ecology and management of riparian shrub communities. USDA Forest Serv. Gen. Tech. Rep. INT-289.
- Bohn, C.C., and J.C. Buckhouse. 1985a. Some responses of riparian soils to grazing management in northeastern Oregon. *J. Range Manage.* 38:378-381.
- Bohn, C.C., and J.C. Buckhouse. 1985b. Coliforms as an indicator of water quality in wildland streams. *J. Soil and Water Cons.* 40:95-97.
- Bryan, K. 1925. Date of channel trenching in the arid Southwest. *Science* 62:338-344.
- Buckhouse, J.C., and G.F. Gifford. 1976. Water quality implications of cattle grazing on a semiarid watershed in southeastern Utah. *J. Range Manage.* 29:109-113.
- Burton, T.A., and S.J. Kozel. 1996. Livestock grazing relationships with fisheries. p. 140-145. In: W.D. Edge, S.L. Olson-Edge (eds.), Sustaining rangeland ecosystems. Oregon State Univ. Extension Service, Special Rep. 953, Corvallis, OR.
- Case, R.L. and J.B. Kauffman. 1997. Wild ungulate influences on the recovery of willows, black cottonwood and thin-leaf alder following cessation of cattle grazing in northeastern Oregon. *Northwest Sci.* 71:115-126.
- Chaney, E., W. Elmore, and W.S. Platts. 1990. Livestock grazing on western riparian areas. Northwest Resource Information Center, Inc. Eagle, Idaho.
- Chaney, E., W. Elmore, and W.S. Platts. 1993. Managing Change: livestock grazing on western riparian areas. Northwest Resource Information Center, Inc. Eagle, Idaho.
- Chapman, D.W., and E. Knudsen. 1980. Channelization and livestock impacts in salmonid habitat and biomass in western Washington. *Trans. Am. Fish. Soc.* 109.
- Claire, E.W., and R.L. Storch. 1977. Streamside management and livestock grazing in the Blue Mountains of Oregon: a case study. p. 111-128, In: Proc. of the workshop on livestock and wildlife-fisheries relationships in the Great Basin. Univ. California, Agric. Station, Sci. Spec. Publ. 3301, Berkeley, CA.
- Clary, W.P. 1995. Vegetation and soil responses to grazing simulation on riparian meadows. *J. Range Manage.* 48:18-25.
- Clary, W.P., E.D. McArthur, D. Bedunah, and C.L. Wambolt (compilers). 1992. Proceedings-Symposium on ecology and management of riparian shrub communities. USDA Forest Serv. Gen. Tech. Rep. INT-289.

- Clary, W.P., and D.E. Medin. 1990. Differences in vegetation biomass and structure due to cattle grazing in a northern Nevada riparian ecosystem. USDA Forest Serv. Re. Pap. INT-427.
- Clary, W.P., and D.E. Medin. 1992. Vegetation, breeding bird, and small mammal biomass in two high-elevation sagebrush riparian habitats. p. 100-110. In: W.P. Clary, E.D. McArthur, D. Bedunah, and C.L. Wambolt (compilers), Proceedings-Symposium on ecology and management of riparian shrub communities. USDA Forest Serv. Gen. Tech. Rep. INT-289.
- Clary, W.P., N.L. Shaw, J.G. Dudley, V.A. Saab, J.W. Kinney, and L.C. Smithman. 1996. Response of a depleted sagebrush steppe riparian system to grazing control and woody plantings. USDA Forest Serv. Res.Pap. INT-RP-492.
- Clary, W.P., and B.F. Webster. 1989. Managing grazing of riparian areas in the intermountain region. USDA Forest Serv. Gen. Tech. Rep. INT-263.
- Davis, L., M. Brittingham, L. Garber, and D. Rourke. 1991. Stream bank fencing. Penn State College of Ag. Sci., Extension Circular 397. University Park, PA.
- Duce, J.T. 1918. The effect of cattle on the erosion of canyon bottoms. *Science* 47:450-452.
- Dudley, T., and M. Embury. 1995. Non-indigenous species in wilderness areas: the status and impacts of livestock and game species in designated wilderness in California Pacific Institute for SIDES, Oakland, CA.
- Duff, D.A. 1977. Livestock grazing impacts on aquatic habitat in Big Creek, Utah. p. 129-142. In: Proc. of the workshop on wildlife-fisheries relationships in the Great Basin. Univ. California, Agric. Station, Sci. Spec. Publ. 3301, Berkeley, CA.
- Elmore, W. 1996. Riparian areas: perceptions in management. USDA Forest Serv., Pacific Northwest Research Station, Natural Resource News 6(3):9.
- Elmore, W., and R.L. Beschta. 1987. Riparian areas: perceptions in management. *Rangelands* 9:260-265.
- Elmore, W., and B. Kauffman. 1994. Riparian and watershed systems: degradation and restoration. p. 212-231. In: M. Vavra, W.A. Laycock, and R.D. Pieper (eds.), Ecological implications of livestock herbivory in the West. Soc. Range Management, Denver, CO.
- Erman, N.A. 1996. Status of aquatic invertebrates. p. 987-1008. In: Sierra Nevada ecosystem project: final report to Congress, Vol. II. Univ. of California, Davis, Centers for Water and Wildlife Resources, Davis, CA.

- Flather, C.H., L.A. Joyce, and C.A. Bloomgarden. 1994. Species endangerment patterns in the United States. USDA Forest Serv. Gen. Tech. Rep. RM-241.
- Fleischner, T.L. 1994. Ecological costs of livestock grazing in western North America. *Cons. Biol.* 8:629-644.
- Gary, H.L., S.R. Johnson, and S.L. Ponce. 1983. Cattle grazing impact on surface water quality in a Colorado front Range stream. *J. Soil Water Cons.* 38:124-128.
- George, M.R. 1996. Creating awareness of clean water issues among private landowners. p. 96-100. In: W.D. Edge, S.L. Olson-Edge (eds.), *Sustaining rangeland ecosystems*. Oregon State Univ. Extension Service, Special Rep. 953, Corvallis, OR.
- Gifford, G.F., and R.H. Hawkins. 1978. Hydrologic impact of grazing on infiltration: a critical review. *Water Resource Res.* 14:305-313.
- Green, D.M., and J.B. Kauffman. 1995. Succession and livestock grazing in a northeast Oregon riparian ecosystem. *J. Range Manage.* 48:307-313.
- Gregory, S.V., F.J. Swanson, W.A. McKee, and K.W. Cummins. 1991. An ecosystem perspective of riparian zones. *BioScience* 41:540-551.
- Gresswell, R.E., B.A. Barton, J.L. Kershner (eds.). 1989. *Practical approaches to riparian resource management*. U.S. Bureau of Land Management, P.O. Box 36800, Billings, Montana.
- Gunderson, D.R. 1968. Floodplain use related to stream morphology and fish populations. *J. Wildl. Manage.* 32:507-514.
- Haveren, B.P., E.B. Janes, and W.L. Jackson. 1985. Nonpoint pollution control on public lands. *J. Soil and Water Cons.* 40(1):92-94.
- Hofmann and R.E. Ries. 1991. Relationship of soil and plant characteristics to erosion and runoff on pasture and range. *J. Soil and Water Cons.* 46(2):143-147.
- Horning, J. 1994. *Grazing to extinction: endangered, threatened and candidate species imperiled by livestock grazing on western public lands*. National Wildlife Federation, Washington, D.C.
- Hubbard, J.P. 1977. Importance of riparian ecosystems: biotic considerations. p. 14-18. In: R.R. Johnson, D.A. Jones, (tech. coords.), *Importance, preservation and management of riparian habitat: A symposium*. USDA Forest Serv. Gen. Tech. Rep. RM-43.
- Hubert, W.A., R.P. Lanka, T.A. Wesche, and F. Stabler. 1985. Grazing management influences on two brook trout streams in Wyoming. p. 290-293. In: R.R. Johnson, C.D. Ziebell, D.R. Patton, and others (tech. coords.), *Riparian ecosystems and their*

management: reconciling conflicting uses. USDA Forest Serv. Gen. Tech. Rep. RM-120.

- Jennings, M.R. 1996. Status of amphibians. p. 921-944. In: Sierra Nevada ecosystem project: final report to Congress, Vol. II. Univ. of California, Davis, Centers for Water and Wildlife Resources, Davis, CA.
- Johnson, R.R., C.D. Ziebell, D.R. Patton, and others (tech. coords.). 1985. Riparian ecosystems and their management: reconciling conflicting uses. USDA Forest Serv. Gen. Tech. Rep. RM-120.
- Johnson, S.R., H.L. Gary, and S.L. Ponce. 1978. Range cattle impacts on stream water quality in the Colorado Front Range. USDA Forest. Serv. Research Note, RM-359.
- Jones, K.B. 1981. Effects of grazing on lizard abundance and diversity in western Arizona. *Southw. Nat.* 26:107-115.
- Kattelman, R., and M. Embury. 1996. Riparian areas and wetlands. p. 201-269. In: Sierra Nevada ecosystem project: final report to Congress, Vol. III. Univ. of California, Davis, Centers for Water and Wildlife Resources, Davis, CA.
- Kauffman, J.B., and W.C. Krueger. 1984. Livestock impacts on riparian ecosystems and streamside management implications...a review. *J. Range Manage.* 37:430-437.
- Kauffman, J.B., W.C. Krueger, and M. Vavra. 1983a. Effects of late season cattle grazing on riparian plant communities. *J. Range Manage.* 36:685-691.
- Kauffman, J.B., W.C. Krueger, and M. Vavra. 1983b. Impacts of cattle on streambanks in northeastern Oregon. *J. Range Manage.* 36:683-685.
- Kleinfelder, D., S. Swanson, G. Norris, and W. Clary. 1992. Unconfined compressive strength of some streambank soils with herbaceous roots. *Soil Sci. Soc. of Amer. Journal* 56:1920-1925.
- Knapp, R.A., and K.R. Matthews. 1996. Livestock grazing, golden trout, and streams in the Golden Trout Wilderness, California: impacts and management implications. *N. Amer. J Fisheries Manage.* 16:805-820.
- Knight, A.W., and R.L. Bottorff. 1984. The importance of riparian vegetation to stream ecosystems. p. 160-167. In: R.E. Warner, K.M. Hendrix (eds), *California riparian systems, ecology, conservation, and productive management.* Univ. of California Press, Berkeley, CA.
- Knopf, F.L., and F.B. Samson. 1994. Scale perspectives on avian diversity in western riparian ecosystems. *Cons. Biol.* 8:669-676.

- Knopf, F.L., J.A. Sedgwick, and R.W. Cannon. 1988. Guild structure of a riparian avifauna relative to seasonal cattle grazing. *J. Wildl. Manage.* 52:280-290.
- Kondolf, G.M. 1993. Lag in stream channel adjustment to livestock exclosure, White Mountains, California. *Restoration Ecology*, December:226-230.
- Kovalchik, B.L., and W. Elmore. 1992. Effects of cattle grazing systems on willow-dominated plant associations in central Oregon. p. 111-119. In: W.P. Clary, E.D. McArthur, D. Bedunah, and C.L. Wambolt (compilers), *Proceedings-Symposium on ecology and management of riparian shrub communities*. USDA Forest Serv. Gen. Tech. Rep. INT-289.
- Lee, L.C., T.A. Muir, and R.R. Johnson. 1989. Riparian ecosystems as essential habitat for raptors in the American West. p. 15-26. In: B.G. Pendleton (ed.), *Western raptor management symposium and workshop*. Nat. Wildl. Fed., Washington D.C.
- Leopold, A. 1946. Erosion as a menace to the social and economic future of the Southwest. *Journal of Forestry* 44:627-633.
- Li, H.W., G.A. Lamberti, T.N. Pearsons, C.K. Tait, J.L. Li, and J.C. Buckhouse. 1994. Cumulative effects of riparian disturbances along high desert trout streams of the John Day Basin, Oregon. *Trans. Amer. Fish. Soc.* 123:627-640.
- Loft, E.R., J.W. Menke, and J.G. Kie. 1991. Habitat shifts by mule deer: the influence of cattle grazing. *J. Wildl. Manage.* 55:16-26.
- Lusby, G.C. 1970. Hydrologic and biotic effects of grazing versus nongrazing near Grand Junction, Colorado. USGS Prof. Paper 700-B:B232-B236.
- Mack, R.N., and J.N. Thompson. 1982. Evolution in steppe with few large, hooved mammals. *American Naturalist* 119:757-772.
- Maloney, S.B., A.R. Tiedemann, D.A. Higgins, T.M. Quigley, and D.B. Marx. 1998. Influence of stream characteristics and grazing intensity on stream temperatures in eastern Oregon. USDA, Forest Serv. Res. Pap. PNW-xxx, (in press).
- Marcuson, P.E. 1977. Overgrazed streambanks depress fishery production in Rock Creek, Montana. p. 143-156. In: *Proc. of the workshop on livestock and wildlife-fisheries relationships in the Great Basin*. Univ. California, Agric. Station, Sci. Spec. Publ. 3301, Berkeley, CA.
- Marlow, C.B., and T.M. Pogacnik. 1985. Time of grazing and cattle-induced damage to streambanks. In: R.R. Johnson, C.D. Ziebell, D.R. Patton, and others (tech. coords.), *Riparian ecosystems and their management: reconciling conflicting uses*. USDA Forest Serv. Gen. Tech. Rep. RM-120.

- Matthews, K.R. 1996. Diel movement and habitat use of California golden trout in the Golden Trout Wilderness. *Trans. Amer. Fisheries Soc.* 125:78-86.
- McInnis, M.L. 1996. Principles of successful livestock grazing in riparian ecosystems. USDA Forest Serv., Pacific Northwest Research Station, *Nat. Res. News* 6(3):1.
- McIntosh, B.A., Sedell, J.R., Smith, J.E., Wissmar, R.C., Clarke, S.E., Reeves, G.H., and L.A. Brown. 1994. Historical changes in fish habitat for select river basins of eastern Oregon and Washington. *Northwest Sci.* 68:36-53.
- Medin, D.E., and W.P. Clary. 1989. Small mammal populations in a grazed and ungrazed riparian habitat in Nevada. USDA Forest Serv. Res. Pap. INT-413.
- Meehan, B.T. 1996. Nature's chisel. *The Oregonian*, February 26, 1996.
- Meehan, W.R. (ed.). 1991. Influences of forest and rangeland management on salmonid fishes and their habitats. American Fisheries Society Special Publ. 19, Bethesda, Maryland.
- Meehan, W.R., and W.S. Platts. 1978. Livestock grazing and the aquatic environment. *J. Soil and Water Cons.* 33:274-278.
- Meehan, W.R., R.J. Swanson, and J.R. Sedell. 1977. Influences of riparian vegetation on aquatic ecosystems with particular reference to salmonid fishes and their food supply. p. 137-145. In: R.R. Johnson, D.A. Jones, (tech. coords.), Importance, preservation and management of riparian habitat: A symposium. USDA Forest Serv. Gen. Tech. Rep. RM-43.
- Myers, T.J., and S. Swanson. 1991. Aquatic habitat condition index, stream type, and livestock bank damage in northern Nevada. *Water Res. Bull.* 27:667-677.
- Myers, T.J., and S. Swanson. 1992. Variation of stream stability with stream type and livestock bank damage in northern Nevada. *Water Res. Bull.* 28:743-754.
- Myers, T.J., and S. Swanson. 1994. Grazing effects on pool forming features in central Nevada. p. 235-244. In: R.A. Marston and V.R. Hasfurther (eds), Effects of human-induced changes on hydrologic systems. Proceedings, Annual Summer Symp. of the Amer. Water Res. Ass., Jackson Hole, Wyoming.
- Myers, T.J., and S. Swanson. 1995. Impact of deferred rotation grazing on stream characteristics in Central Nevada: a case study. *North Amer.J. of Fisheries Manage.* 15:428-439.
- Myers, T.J., and S. Swanson. 1996a. Long-term aquatic habitat restoration: Mahogany Creek, Nevada, as a case study. *J. of the Amer. Water Res. Ass.* 32:241-252.

- Myers, T.J., and S. Swanson. 1996b. Temporal and geomorphic variations of stream stability and morphology: Mahogany Creek, Nevada. *J. of the Amer. Water Res. Ass.* 32:253-265.
- Ohmart, R.D. 1996. Historical and present impacts of livestock grazing on fish and wildlife resources in western riparian habitats. p. 245-279. In: P.R. Krausman (ed.), *Rangeland wildlife. Soc. for Range Manage., Denver CO.*
- Ongerth, J.E., and H.H. Stibbs. 1987. Identification of *Cryptosporidium* oocysts in river water. *Appl. and Environ. Microb.* 53:672-676.
- Oregon Department of Environmental Quality. 1995a. Temperature, 1992-1994 Water Quality Standards Review. Oregon Department of Environmental Quality, 811 Sixth Avenue, Portland OR.
- Oregon Department of Environmental Quality. 1995b. Dissolved Oxygen, 1992-1994 Water Quality Standards Review. Oregon Department of Environmental Quality, 811 Sixth Avenue, Portland OR.
- Orr, H.K. 1975. Recovery from soil compaction on bluegrass range in the Black Hills. *Transactions of the ASAE:* 1076-1081.
- Overton, C.K., G.L. Chandler, J.A. Pisano. 1994. Northern/Intermountain regions' fish habitat inventory: grazed, rested, and ungrazed reference stream reaches, Silver King Creek, California. *USDA For. Serv. Gen. Tech. Rep. INT-GTR-311.*
- Owens, L.B., W.M. Edwards, and R.W. Van Keuren. 1989. Sediment and nutrient losses from an unimproved, all-year grazed watershed. *J. Environ. Qual.* 18:232-238.
- Owens, L.B., W.M. Edwards, and R.W. Van Keuren. 1996. Sediment losses from a pastured watershed before and after stream fencing. *J. Soil and Water Cons.* 51:90-94.
- Peterman, R.M. 1990. Statistical power analysis can improve fisheries research and management. *Can. J. Fish. Aquat. Sci.* 47:2-15.
- Platts, W.S. 1981a. Sheep and streams. *Rangelands* 3:158-160.
- Platts, W.S. 1981b. Influence of forest and rangeland management on anadromous fish habitat in western North America: 7. Effects of livestock grazing. *USDA Forest Serv. Gen. Tech. Rep. PNW-124.*
- Platts, W.S. 1982. Livestock and riparian-fishery interactions: what are the facts? *Trans. North Amer. Wildl. and Nat. Res. Conf.* 47:507-515.

- Platts, W.S. 1991. Livestock grazing. p. 389-424. In: W.R. Meehan (ed.), Influences of forest and rangeland management on salmonid fishes and their habitats. Amer. Fisheries Soc. Sp. Publ. 19:389-423.
- Platts, W.S., and F.J. Wagstaff. 1984. Fencing to control livestock grazing on riparian habitats along streams: is it a viable alternative? N. Amer. J. Fish. Manage. 4:266-272.
- Ponce, V.M., and D.S. Lindquist. 1990. Management of baseflow augmentation: a review. Water Res. Bull. 26:256-268.
- Popolizio, C.A., H. Goetz, and P.L. Chapman. 1994. Short-term response of riparian vegetation to 4 grazing treatments. J. Range Manage. 47:48-53.
- Reiser, D.W., and T.C. Bjornn. 1979. Influence of forest and rangeland management on anadromous fish habitat in the western United States and Canada, 1. Habitat requirements of anadromous salmonids. USDA Forest Serv. Gen. Tech. Rep. PNW-96.
- Rhodes, J.J., McCullough, D.A., and F.A. Espinosa, Jr. 1994. A coarse screening process for evaluation of the effects of land management activities on salmon spawning and rearing habitat in ESA consultations. Columbia River Intertribal Fish Comm. Tech. Rept. 94-4, Portland, OR.
- Rinne, J.N. 1985. Livestock grazing effects on southwestern streams: a complex research problem. p. 295-300. In: R.R. Johnson, C.D. Ziebell, D.R. Patton, and others (tech. coords.), Riparian ecosystems and their management: reconciling conflicting uses. USDA Forest Serv. Gen. Tech. Rep. RM-120.
- Rinne, J.N. 1988. Effects of livestock grazing exclosure on aquatic macroinvertebrates in a montane stream, New Mexico. Great Basin Nat. 48:146-153.
- Roath, L.R., and W.C. Krueger. 1982. Cattle grazing and influence on a forested range. J. of Range Manage. 35:332-338.
- Rostagno, C.M. 1989. Infiltration and sediment production as affected by soil surface conditions in a shrubland of Patagonia, Argentina. J. Range Manage. 42:382-385.
- Saab, V.A., C.E. Bock, T.D. Rich, and D.S. Dobkin. 1995. Livestock grazing effects in western North America. p. 311-353. In: T.E. Martin, D.M. Finch (eds.), Ecology and management of neotropical migratory birds: a synthesis and review of critical issues. Oxford University Press, London.
- Samson, F.B., F.L. Knopf, and L.B. Hass. 1988. Small mammal response to the introduction of cattle into a cottonwood floodplain. p. 432-438. In: R.C. Szaro, K.E. Severson, D.R. Patton, (tech. coords.), Management of amphibians, reptiles, and small mammals in North America. USDA For. Serv. Gen. Tech. Rep. RM-166.

- Sarr, D., R.A. Knapp, J. Owens, T. Balsler, and T. Dudley. 1996. Ecosystem recovery from livestock grazing in the southern Sierra Nevada. Aldo Leopold Wilderness. Res. Insti., Missoula, MT.
- Savory, S. 1988. Holistic Resource Management. Island Press, Covalo, CA.
- Schepers, J.S., and D.D. Francis. 1982. Chemical water quality of runoff from grazing land in Nebraska: I. Influence of grazing livestock. *J. Environ. Qual.* 11:351-354.
- Schepers, J.S., B.L. Hackes, and D.D. Francis. 1982. Chemical water quality of runoff from grazing land in Nebraska: II. Contributing factors. *J. Environ. Qual.*, 11:355-359.
- Schulz, T.T., and W.C. Leininger. 1990. Differences in riparian vegetation structure between grazed areas and exclosures. *J. Range Manage.* 43:295-299.
- Schulz, T.T., and W.C. Leininger. 1991. Nongame wildlife communities in grazed and ungrazed riparian sites. *Great Basin Nat.* 51:286-292.
- Sedgwick, J.A., and F.L. Knopf. 1987. Breeding bird response to cattle grazing of a cottonwood bottomland. *J. Wildl. Manage.* 51:230-237.
- Sedgwick, J.A., and F.L. Knopf. 1991. Prescribed grazing as a secondary impact in a western riparian floodplain. *J. Range Manage.* 44:369-373.
- Shaw, N.L. 1992. Recruitment and growth of Pacific willow and sandbar willow seedlings in response to season and intensity of cattle grazing. p. 130-137, In: W.P. Clary, E.D. McArthur, D. Bedunah, and C.L. Wambolt (compilers), Proceedings-Symposium on ecology and management of riparian shrub communities. USDA Forest Serv. Gen. Tech. Rep. INT-289.
- Skovlin, J.M. 1984. Impacts of grazing on wetlands and riparian habitat: a review of our knowledge. p. 1001-1103. In: *Developing strategies for range management.* Westview Press, Boulder, CO.
- Stacey, P.B. 1995. Diversity of rangeland bird populations. p.33-41. In: N.E. West (ed.), *Biodiversity on rangelands.* College of Natural Resources, Utah State University, Logan, UT.
- Stephenson, G.R., and R.C. Rychert. 1982. Bottom sediment: a reservoir of *Escherichia coli* in rangeland streams. *J. Range Manage.* 35:119-123.
- Stephenson, G.R., and L.V. Street. 1978. Bacterial variations in streams from a southwest Idaho rangeland watershed. *J. Environ. Qual.* 7:150-157.
- Stevens, R., E.D. McArthur, and J.N. Davis. 1992. Reevaluation of vegetative cover changes, erosion, and sedimentation on two watersheds--1912-1983. p. 123-128. In:

W.P. Clary, E.D. McArthur, D. Bedunah, and C.L. Wambolt (compilers),
Proceedings-Symposium on ecology and management of riparian shrub communities.
USDA Forest Serv. Gen. Tech. Rep. INT-289.

Stoddart, L.A., and A. Smith. 1955. Range management, 2nd edition. McGraw-Hill, New York, NY.

Stuber, R.J. 1985. Trout habitat, abundance, and fishing opportunities in fenced vs. unfenced riparian habitat along sheep creek, Colorado. p. 310-314. In: R.R. Johnson, C.D. Ziebell, D.R. Patton, and others (tech. coords.), Riparian ecosystems and their management: reconciling conflicting uses. USDA Forest Serv. Gen. Tech. Rep. RM-120.

Szaro, R.C. 1989. Riparian forest and scrubland community types of Arizona and New Mexico. *Desert Plants* 9(3-4):72-138.

Szaro, R.C., S.C. Belfit, J.K. Aitkin, and J.N. Rinne. 1985. Impacts of grazing on a riparian garter snake. p. 359-363. In: R.R. Johnson, C.D. Ziebell, D.R. Patton, and others (tech. coords.), Riparian ecosystems and their management: reconciling conflicting uses. USDA Forest Serv. Gen. Tech. Rep. RM-120.

Tait, C.K., J.L. Li, G.A. Lamberti, T.N. Pearsons, and H.W. Li. 1994. Relationships between riparian cover and community structure of high desert streams. *J. N. A. Benthol. Soc.* 13:45-56.

Taylor, D.M. 1986. Effects of cattle grazing on passerine birds nesting in riparian habitat. *J. Range Manage.* 39:254-258.

Taylor, F.R., L.A. Gillman, and J.W. Pedretti. 1989. Impact of cattle on two isolated fish populations in Pahrnagat Valley, Nevada. *Great Basin Nat.* 49:491-495.

Thurrow, T.L. 1991. Hydrology and erosion. p.141-159. In: R.K. Heitschmidt, and J.W. Stuth (eds.), *Grazing management: an ecological perspective*. Timber Press, Portland OR.

Thomas, J.W., C. Maser, and J.E. Rodiek. 1979. Wildlife habitats in managed rangelands-- The Great Basin of southeastern Oregon: riparian zones. USDA Forest Serv. Gen. Tech. Rep. PNW-80.

Tiedemann, A.R., and D.A. Higgins. 1989. Effects of management strategies on water resources. p.56-91. In: T.M. Quigley, H.R. Sanderson, and A.R. Tiedemann, *Managing interior Northwest rangelands: The Oregon Range Evaluation Project*. USDA Forest Serv. Gen. Tech. Rep. PNW-GTR-238.

Tiedemann, A.R., D.A. Higgins, T.M. Quigley, H.R. Sanderson, and D.B. Marx. 1987. Responses of fecal coliform in streamwater to four grazing strategies. *J. Range Manage.* 40:322-329.

- Trimble, S.W. 1994. Erosional effects of cattle on streambanks in Tennessee, U.S.A. *Earth Surface Processes and Landforms* 19:451-464.
- Trimble, S.W., and A.C. Mendel. 1995. The cow as a geomorphic agent -- a critical review. *Geomorphology* 13:233-253.
- U.S. Department of Interior. 1993. Riparian area management, process for assessing proper functioning condition. TR 1737-9 1993, Bureau of Land Management, Box 25047, Denver, CO.
- U.S. Department of Interior. 1994a. Rangeland reform '94, Draft environmental impact statement. Bureau of Land Management, Washington, D.C.
- U.S. Department of Interior. 1994b. Western riparian wetlands (Chapter 12). p. 213-238. In: *The impact of federal programs on wetlands, Vol. II, A report to Congress by the Secretary of the Interior*, Washington D.C., U.S. Fish and Wildlife Service, Arlington, VA.
- U.S. Environmental Protection Agency. 1995. National Water Quality Inventory, 1994 Report to Congress Executive Summary. Office of Water, Washington DC 20460.
- U.S. General Accounting Office. 1988. Public rangelands: some riparian areas restored by widespread improvement will be slow. GAO/RCED-88-105.
- Warner, R.E., and K.M. Hendrix (eds). 1984. *California riparian systems, ecology, conservation, and productive management*. Univ. of California Press, Berkeley, CA.
- Weller, M.W. 1996. Birds of rangeland wetlands. p. 71-82. In: P.R. Krausman (ed.), *Rangeland wildlife*. The Society of Range Management, Denver CO. White, R.J., and O.M. Brynildson. 1967. Guidelines for management of trout stream habitat in Wisconsin. Dep. Nat. Resour. Tech. Bull. 39:, Madison, WI.
- Winegar, H.H. 1977. Camp Creek channel fencing -- plant, wildlife, soil, and water response. *Rangeman's J.* 4:10-12.