Allotment Management Plan Williamson Valley Allotment

USDA Forest Service Chino Valley Ranger District, Prescott National Forest Yavapai County, Arizona

Prepared by: John A. Kava, Rangeland Management Specialist	Date:	11/17
Reviewed by/ agreed to May Harmittee	Date_2	1/17
Approved By Omero Torres, Chino Valley District Ranger	Date_2	4/17

Introduction

This Allotment Management Plan (AMP) is a direct result of the Environmental Assessment and subsequent Decision Notice (DN), dated September 30, 2016.

The Williamson Valley Allotment is located on the Chino Valley District of the Prescott National Forest (PNF) and represents the project area for this analysis, an area of approximately 49,000 acres. The allotment is located in the southwest portion of the district, approximately 16 miles west of Chino Valley, Arizona

Roughly two thirds of the allotment is in the Big Chino Watershed and one third in the Santa Maria River watershed. Elevation ranges from 4,750 feet in Hitt Wash and Williamson Valley Wash on the eastern boundary to 7,200 feet on Camp Wood Mountain in the northwest corner of the allotment. The topography is rough and broken on most of the allotment with more gentle hills around Hitt and Williamson Valley Washes on the northeastern section.

Vegetation on the allotment consists primarily of Pinyon Juniper evergreen shrub and Interior Chaparral. Canopy cover from shrub species is moderately to extremely thick in some locations to the extent that herbaceous forage is reduced or absent. A portion of the forage base of the allotment is provided by desirable browse species such as turbinella oak with mountain mahogany, deerbrush, and skunkbush found in smaller quantities. Perennial grasses can be locally abundant, especially in juniper woodlands that have been previously thinned, and south aspects. Important forage grasses on the allotment include blue grama, sideoats grama, threeawns, sand dropseed, and squirreltail.

The northeastern two thirds of the allotment is drained by Pine Creek, Hitt Wash, and Williamson Valley Wash which are tributaries of the upper Verde River. Riparian vegetation occurs along these stretches. Hitt Wash and Williamson Valley Wash have mainly herbaceous riparian vegetation such as sedges, rushes, horsetails, and other grass-like plants. There are localized areas of velvet ash, willow and cottonwood.

Desired Conditions

The desired conditions on this grazing allotment, based on the Forest Plan (see Attachment 1 Forest Plan Standards and Guidelines) and the work of the Interdisciplinary Analysis Team (ID team), include:

- Range administration that provides for the maintenance of satisfactory Rangeland Management Status (RMS) with a static or upward apparent trend;
- The maintenance of vegetation with mid- to high similarity to the Desired Vegetative Status (DVS) providing for ecological functionality and resiliency following disturbance while sustaining long-term productivity of the land;
- The installation and maintenance of structural improvements, such as water-supply systems, that enhance management control and flexibility and allow for effective distribution of forage use;
- The maintenance of soils in satisfactory condition over the long-term, or show improvement in areas departing from satisfactory condition where livestock grazing is contributing to the departure;



- The maintenance of functioning spring-fed riparian systems, and saturated soils where
 potential exists, that support vegetation within site potential and provide habitat for
 riparian-dependent plants and animals while providing water sources for wildlife and
 livestock needs;
- The maintenance of fully functional riparian systems supported by herbaceous and multiage woody vegetation, within site potential, that provides for stable stream channels and banks and habitat for riparian-dependent plants and animals;
- · Protection and preservation of important historic and cultural sites; and
- The maintenance of suitable habitats for Management Indicator Species, Migratory Bird Treaty Act species, Forest Service Sensitive species, and for indigenous plant and animal species.

Resource Objectives

The following management objectives were developed to measure progress towards meeting desired conditions:

- Improve or maintain cover of perennial grasses to achieve mid- to high similarity with the
 potential perennial grass canopy cover and composition as shown in the Ecological
 Classification for the Prescott National Forest for key TEUI map units; achieve an
 upward trend in vegetation condition towards this objective.
 - In Little Pine, Burnt and Whiskey pastures, TEUI 48; Tailholt pasture, TEUI 490;
 and Camp Wood pasture, TEUI 542; detect an improvement of vegetation spatial distribution (GAP) within 5-7 years (soil health).
 - Humphreys Pasture (East): TEUIs 461, 462, 481, and 490 are in unsatisfactory soil condition. The management objective is to not exacerbate soil damage through livestock use.

Grazing Management

Permitted Numbers, Season of Use, and Animal Unit Months

# of Livestock	Season of Use	Animal Unit Months
Between 225 to 300 head of cattle, cow/calf pairs and bulls	yearlong	Not to exceed 3,600 Animal- Unit- Months ¹

The period of grazing and the stocking numbers on NFS lands will be determined by monitoring, designated in the Annual Operating Instructions (AOI) and authorized in the Bill for Collection.

AOI will be prepared each year in cooperation with the permittee to allow for consideration of current allotment conditions and management objectives. This AOI will detail the current season's grazing schedule, the stocking level, the improvement maintenance needs, needed improvements, and the allowable use levels on key forage and browse species.

Annual stocking would be based on adaptive management, considering forage

¹ Animal-Unit-Month (AUM) is the amount of oven-dry forage required by one mature cow of about 1,000 pounds, either dry or with a calf up to six months of age, or their equivalent, for a standardized period of 30 animal-unit-days.



production, water availability, and resource conditions. Annual stocking could fall below the low end of the proposed stocking range. There are six larger pastures and five smaller pastures used in a rotational grazing system. Pasture rest and deferment will be scheduled to provide for achieving desired resource conditions.

Adaptive management is designed to provide sufficient flexibility to allow livestock management to address changes in climatic conditions, seasonal fluctuations in forage production, and other dynamic influences on the ecosystem in order to effectively make progress toward or maintain desired conditions of the rangeland and other resources. Under the adaptive management approach, regular/annual monitoring of short-term indicators may suggest the need for administrative changes in livestock management. If monitoring indicates that progress toward desired conditions is not being achieved on the allotment, management will be modified.

Modifications can include adjustments in timing, intensity, and duration of grazing. Timing is the time of year the livestock are present in a pasture. Intensity is the degree to which forage is removed through grazing and trampling by livestock. Duration is the length of time livestock are present in a given pasture. These modifications would be made through administrative decisions such as: the specific number of head stocked on the allotment annually or in a particular season; the class of animals stocked (cow/calf pairs vs. yearlings, steers or heifers, etc.); specific dates of grazing; livestock herd movement; and periods of rest, deferment, or non-use of portions or all of the allotment for an appropriate period of time, as conditions warrant. Such changes will not result in exceeding the AUMs authorized for livestock use.

Application of standard management practices such as salting, herding, and controlling access to water to achieve proper distribution or lessen the impact on areas which are sensitive or are natural concentration areas will be applied by the permittee.

Protein, salt, and other supplements will not be placed within ¼ mile of water or any identified sensitive plant population. New improvements (e.g. pipelines, troughs, tanks, or fences) will be designed to avoid adverse impacts to any such populations.

Allowable Use

Allotment Wide Measures:

Grazing intensity guidelines will be applied across the allotment to provide rangeland managers with information needed to adapt management through adjustments, as may be needed, on an annual basis. Examples of appropriate grazing intensity and forage use guidelines for areas of the allotment that are generally described to be in satisfactory condition include:

- A management guideline of 35-45% utilization of key forage plants in upland key areas as measured at the end of the growing season or seasonal use period;
- Up to 50-60% leaders browsed on key upland woody species;
- Minimum stubble height on key riparian herbaceous species: four to six inches where sedges and rushes are key and eight inches where deergrass is key;
- Up to 20% use by weight on key woody species within riparian areas; or less than 50% of terminal leaders browsed on woody species less than 6 feet tall.

Grazing intensity will be determined using key herbaceous and browse species within key areas. Grazing would be adjusted if periodic monitoring indicates that desired resource conditions are not being maintained.



Site-specific Measures:

The following measures will be applied in areas of concern where current conditions are not meeting desired conditions, and management objectives have been established to measure progress towards meeting desired resource conditions:

- Light utilization levels (30% or less) and integrated rest periods for <u>Little Pine</u> and <u>Whiskey</u> pastures (TEUI 48), <u>Tailholt</u> pasture (TEUI 490).
- Incidental (0-30%) grazing <u>Humphreys</u> pasture (east) (TEUIs 461, 462, 481, and 490). No practices to draw livestock (water and supplement placement) would occur.
- Rest or deferment,
 - <u>Burnt</u> pasture (TEUI 48), to allow freeze-thaw cycles to improve soil compaction.
 - <u>Camp Wood</u> pasture (TEUI 542), to encourage grass plant establishment and litter development that can alleviate soil compaction.

Once desired conditions for vegetation or soil are being met in areas needing improvement, then the allotment-wide utilization standards could be applied.

In the event that the above resource protection measures do not accomplish site-specific resource objectives, additional optional measures may be implemented. These optional measures will be designed to address site-specific resource concerns and may include, but are not limited to, such things as temporary fencing, electric fencing, and reconstruction of existing non-functional improvements and construction of new improvements such as drift fences.

Rangeland Improvement Program

Construction of New Range Improvements:

Construction of the following new structural improvements has been approved to address resource concerns. These improvements are intended to aid in the achievement or maintenance of desired resource conditions by improving livestock distribution. The Forest Service will work in collaboration with the Permittee, AZ G&F and other partners as the opportunity presents.

Different types of water developments may be employed depending on the location, and could include trick tanks with a pipeline to water troughs, earthen stock tanks, or wells.

Construct 12 additional water sources in the following locations:

- Upper Hitt Pasture, section 15 (likely a trick tank), and SE quarter of section 16 on the pasture division fence;
- Tailholt Pasture, SE quarter of section 22;
- Lower Hitt Pasture, SW quarter of section 25;
- Shared water source between Burnt and Upper Hitt Pastures in NW quarter of section 26;
- Burnt Pasture south half of section 27;
- Shared water source for Whiskey and Brushy Pastures in SW quarter of section 33;
- Brushy Pasture NW quarter of section 36;
- Stinson Pasture
 - SE quarter of section 17,

Page // of 27

- o SW quarter of section 29,
- SW quarter of section 31;
- Camp Wood pasture north half of section 33.

Convert 3 existing earthen stock tanks to trick tanks in order to provide more reliable water supplies:

- Cottonwood Pasture,
 - Section 23 Tank,
 - o Coldwater Tank
- Tailholt Pasture, tank in SW quarter of section 10.

Construct a new holding pasture south of Spades Tank in the Tailholt Pasture.

Maintenance Responsibility

The Term Grazing Permit includes a list of all improvements which the permittee will continue to maintain at a level that effectively provides for their intended uses and purposes. Range improvements will be inspected periodically during the term of the permit to document condition.

Damage resulting from big game, wind, other acts of nature, or human caused actions, must be repaired in a timely manner so as to ensure the integrity of the structures.

All maintenance of exterior fences must be completed prior to turn-on each year. (It is the responsibility of the permittee to ensure that the necessary coordination occurs between adjacent allotments to ensure maintenance is completed in a timely manner).

AOI will identify range improvements in need of maintenance. Existing improvements may be replaced when their conditions warrant. All improvements identified on allotment maps have been evaluated and determined necessary to the management of the allotment through the life of this plan.

Access to Improvements:

Authorization for cross-country motorized travel is provided for the permittee to administer the livestock operation and maintain improvements under the terms and conditions of the Term Grazing Permit.

Annual authorization for actions implementing management direction in the AMP will be included in the AOI, such as a description of the anticipated level of cross-county travel, travel needed for improvement maintenance, new improvement construction, or reconstruction of existing improvements.

All authorizations for cross-country motorized travel are subject to existing regulations intended to protect natural and/or heritage resources. Cross-country travel is not allowed when such travel would cause unacceptable resource damage.

The permittee may be authorized to maintain forest systems road to facilitate the maintenance, replacement, or installation of range improvements. Maintenance activities will adhere to Forest Service standards and be authorized through a road use permit.

Page 12 of 27

Drought Management

Perennial grasses and major browse species need deferment/rest in order to provide time to recover from drought induced stress. Even when rested or deferred, if adequate precipitation is not received, recovery may not be adequate for livestock use.

Move cattle to the next scheduled pasture when utilization in pastures is met. If complete removal of livestock is necessary, they may be authorized to return to the allotment once conditions improve; meaning sufficient recovery from the effects of drought stress has occurred and there has been enough herbaceous production to support livestock numbers. Potential return of livestock will be evaluated no earlier than the summer growing season.

Monitoring and Evaluation

Implementation Monitoring

This monitoring will be conducted on an annual basis and will include such things as livestock actual use (# of head, # of months) and scheduled and unscheduled inspections to ensure that all livestock and grazing management measures stipulated in the permit, AMP, and AOI are being implemented (e.g. cattle numbers, on/off dates, rotation schedules, maintenance of improvements, mitigation measures).

Monitoring activities would be focused on those resources that need improvement or where there is a concern for an important habitat type. For this project, there are soil and vegetation condition concerns in the Little Pine, Burnt, Whiskey pastures in TEUI 48, Tailholt pasture TEUI 490, and Camp Wood Pasture TEUI 452.

 Canopy gap and vegetative ground cover will be measured at key areas in Little Pine, Burnt and Whiskey pastures (TEUI 48), Tailholt pasture (TEUI 490), Camp Wood pasture (TEUI 542) Results of monitoring will be analyzed against baseline data or Ecological Classification description to determine if objectives are being met.

Forage utilization will be monitored on the allotment at key areas and at areas identified with site-specific resource concerns. See Attachment 2, Key Area Map.

The key area concept is based on the premise that no range of appreciable size will be grazed uniformly (Holechek, Pieper and Herbel, 1998). When key areas are "properly" used there may be substantial areas that are used more or less than the key areas, including some that will not be used at all. Forest Service personnel can work with the permittee in selecting these areas.

(Monitoring of allowable use on key forage species in key areas is the joint responsibility of the Forest Service and the permittee. Although the Forest Service will make every effort to assist the permittee in ensuring compliance with standards, the permittee has the ultimate responsibility for ensuring that the allowable use standards are met).

If periodic field checks indicate that plant vigor or production is poor, and bare soil is increasing, this would trigger a need to make adaptive management adjustments. This could also result in reevaluation of vegetation or soil condition through effectiveness monitoring.

Field Checks will include informal inspections, formal inspections, and permittee compliance monitoring.

Informal Inspections

Page 13 of 27

Informal inspections conducted by the Forest Officer will be made as the opportunity arises, such as when the Forest Officer is working in the area or is passing through the allotment.

The permittee will be notified by telephone of any significant observations needing immediate attention. Significant observations will be documented in writing by the Forest Officer and a copy of the inspection notes will be sent to the permittee in a timely manner.

Formal Inspections

Formal inspections conducted by the Forest Officer will be made as time and competing duties allow with an attempt to inspect each of the pastures.

The permittee will be requested to accompany the Forest Officer during the inspections. Significant findings from these inspections will be documented in a letter or inspection report sent to the permittee in a timely manner.

Permittee Compliance Monitoring

The permittee will:

- Monitor the allotment continuously throughout the grazing season to determine current resource conditions and to ensure the terms of the permit are being met.
- Document all findings through notes, photographs, or other means decipherable by the Forest Officer
- o Share monitoring information with the Forest Officer, and
- o Coordinate with the Forest Officer to resolve any problems that arise.

Effectiveness Monitoring

The permittee is encouraged to participate in any effectiveness (e.g. long term condition and trend) monitoring and evaluation conducted on the allotment. This type of monitoring evaluates the success of management in achieving the desired objectives within key and critical areas or on permanent transects at an interval of 10 years or less. Data collected for the Allotment Management Plan revision serves as a baseline for vegetation and soil condition. The same key areas evaluated for the analysis will be re-visited to determine if desired conditions are being maintained, or there is acceptable progress in those areas needing improvement. Effectiveness monitoring may also be conducted if data and observations from implementation monitoring indicate a need.

Both qualitative and quantitative monitoring methods will be used in accordance with Interagency Technical References, the Region 3 Rangeland Analysis and Management Training Guide, and the Region 3 Allotment Analysis Handbook. Common methods to evaluate vegetation trend include plant frequency changes over time, or comparison of existing vegetation canopy cover and species composition to the potential natural vegetation based on soil type, climate, elevation, topography, and past land uses.

Page 14 of 27

Attachment 1, Relevant Forest Plan Standards and Guidelines

Range Management		
Std- Range-1	Water troughs shall incorporate escape devices to prevent animal entrapments	
Std- Range-2	Year-long livestock grazing in riparian areas (streams, springs, and seeps) shall be avoided to prevent adverse impacts to water quality and riparian habitat in those areas	
Guide- Range-1	The placement of salt, minerals, and/or other supplements for the purposes of livestock management should be located further than one-quarter mile from riparian areas or seasonally present water.	
Guide- Range-2	 For structural improvements: Implement design features that incorporate wildlife needs and reduce barriers to movement and entrapment hazards Consider wildlife needs in fence placement and design to reduce barriers and hazards to movement and minimize chances of entrapment Remove fencing when it is no longer needed 	
Guide- Range-3	After occurrence of wildland fire or mechanical activity that removes most vegetation, a time period for recovery, establishment, and regrowth of vegetation should be determined and applied to meet site-specific objectives	
Guide- Range-4	Livestock salting should be located away from known locations of Southwestern Region sensitive plant species so that plants are not adversely affected by associated trampling	
Guide- Range-5	Livestock use of woody riparian species (e.g. cottonwood, willow, ash, and alder) should provide for maintenance of those species and allow regeneration of new individuals leading to diverse age classes of woody riparian species where potential for native woody vegetation exists	
Guide- Range-6	Grazing intensity, frequency, occurrence, and period should provide for growth and reproduction of desired plant species while maintaining or enhancing habitat for wildlife	
	Watersheds guidelines	
Guide- WS-4	Adverse impact to stream channel features (e.g. streambanks, obligate riparian vegetation) should be minimized by modifying management actions. Examples of modification could include, but are not limited to: adjusting timing and season of grazing, limiting use and location of heavy machinery, or avoiding placing trails or other recreation structures where recreation use could negatively affect stream channel features	
Guide- WS-5	Ground cover sufficient to filter runoff and prevent erosion should be retained in riparian corridors, seeps, and springs	
Guide- WS-9	Along perennial streams, perennial intermittent streams, and spring ponds, mitigation such as offsite water for livestock should be provided to reduce impacts on riparian communities and groundwater dependent sites	

Guide-WS- 10	Measures that restrict use should be considered as a way to mitigate recurring negative impacts to aquatic species and riparian plants. These could include, but are not limited to: installation of barriers, road closures, area closures, or seasonal restrictions
	Soils Guideline
Guide- Soil-1	Projects should be designed to limit activities that would cause long term impacts to soils such as loss of ground cover, severely burned soils, detrimental soil displacement, erosion, puddling, or compaction. Where disturbance cannot be avoided, project-specific soil and water conservation practices should be developed.
	Vegetation Standard
STD-Veg-2	When treating nonnative and invasive plant species, design features in appendix B of the "Final Environmental Impact Statement for Integrated Treatment of Noxious or Invasive Weeds" (Forest Service, 2005a) or the most current direction must be followed to protect endangered, threatened, proposed, and candidate wildlife and plant species and their habitats

Attachment 2, Key Area Map Williamson Valley Allotment Legend

Page 11 of 11